



# Rx Cefquinome Sulphate Injection (Vet)

COBACTAN® 2.5%



NOT FOR HUMAN USE, FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG – CAUTION  
Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

COMPOSITION

Cefquinome sulphate equivalent to Cefquinome 25 mg per ml.

TARGET SPECIES

Cattle and pigs

INDICATIONS

For the treatment of infection caused by susceptible organisms in cattle and pigs.

DOSAGE AND ADMINISTRATION

SPECIES	DOSAGE	FREQUENCY
Cattle	1 mg Cefquinome / kg Bodyweight (equivalent to 2 ml / 50 kg bw)	Daily for 3-5 consecutive days
Pigs	2 mg Cefquinome / kg bodyweight (equivalent to 2 ml per 25 kg bw)	Daily for 3 consecutive days

All treatments to be given by intramuscular injection. Studies have indicated the advisability of giving second and subsequent injections at a different injection site.

Shake the vial well before using.

The product does not contain an antimicrobial preservative. Swab the septum before removing each dose. Use a dry sterile needle and syringe. The cap may be safely punctured up to 25 times. When treating groups of animals, use a draw-off needle.

WITHDRAWAL PERIODS

Meat and milk from treated animals may not be used for human consumption during treatment. Meat and milk may not be taken for human consumption until the following times after the final treatment.

SPECIES	PRODUCT	WITHDRAWAL PERIOD
Cattle	Meat	5 days
Cattle	Milk	1 day
Pigs	Meat	3 days

CONTRAINDICATIONS

Hypersensitivity to Cephalosporins is rare; however Cefquinome should not be administered to animals which are known to be hypersensitive to  $\beta$ -lactam antibiotics.  
Do not administer to animals less than 1.25 kg body weight.

ADVERSE EFFECTS

Use of the product may result in localized tissue reaction. Tissue lesions are repaired 15 days after the last administration of the product.

SPECIAL WARNINGS

Penicillins and Cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross sensitivity to Cephalosporins and vice versa.  
Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitized, or if you have been advised not to work with such preparation.
2. Handle this product with great care to avoid exposure, taking all recommended precautions.
3. If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips eyes or difficulty with breathing, are more serious symptoms and require urgent medical attention.

SHELF LIFE: 24 months

SPECIAL STORAGE CONDITIONS

- Store in a refrigerator (+2°C to +8°C) and protect from light.
- Following withdrawal of the first dose, use the remaining content within 4 weeks.
- Discard unused material.
- Keep out of reach of children.

PRESENTATION

50 ml glass vial.

Manufactured by:

Intervet International GmbH  
Feldstrasse 1A,D-85716 Unterschleissheim,  
Germany.

**Imported & Marketed by:** Intervet India Pvt. Ltd.,  
Bldg. No. B-1, Godown No.12 to 16, Ground Floor & Godown No.12-A to 17-A First Floor, House No. 233,  
Sagar Complex, Owali, Bhiwandi, Thane -421302, Maharashtra, India.

**®Registered Trademark**

