


LIVE, FREEZE-DRIED VACCINE AGAINST *BORDETELLA BRONCHISEPTICA* AND CANINE PARAINFLUENZA VIRUS

362294 R1



Nobivac® KC



NOT FOR HUMAN USE; FOR ANIMAL TREATMENT ONLY

Warning: To be sold by retail on the prescription of a Veterinary Doctor only.

Nobivac® KC is a live freeze-dried vaccine containing live *Bordetella bronchiseptica* strain B-C2 and live Canine parainfluenza virus strain Cornell for active immunisation of dogs against *Bordetella bronchiseptica* and Canine parainfluenza virus, resulting in prevention of kennel cough.

Composition:

Each dose of 0.4 ml of reconstituted vaccine contains,
- at least $10^{8.0}$ cfu of live *Bordetella bronchiseptica* strain B-C2 and
- at least $10^{3.0}$ TCID₅₀ of live canine parainfluenza virus strain Cornell

Target Species: Dogs

Indications:

Active immunisation of dogs against *Bordetella bronchiseptica* and Canine parainfluenza virus, resulting in prevention of Kennel cough disease.

Vaccination Scheme:

One dose at least 72 hours prior to period of anticipated risk.
Dogs should be at least two weeks of age.

Revaccination:

Annually.

Dose and administration:

Allow the diluent to reach room temperature. Aseptically reconstitute the freeze-dried vaccine with the sterile diluent provided. Fill 1 ml disposable syringe with 1 dose (0.4 ml) of vaccine, remove the needle, connect the applicator tip and administer 0.4 ml into one nostril. The vaccine should be used within 1 hour after reconstitution.

Contra-Indications and Warning:

- Only healthy dog should be vaccinated and an adequate clinical examination should be performed prior to inoculation.
- The vaccine is safe for pregnant bitches.

Compatibility:

Do not administer in conjunction with other intranasal treatments or during antibiotic treatment.

Immunity:

Onset of immunity:
- for *Bordetella bronchiseptica*: 72 hours after vaccination
- for canine parainfluenza virus: three weeks after vaccination
Duration of immunity: 1 year

Vaccination Reaction:

Mild discharge from the eyes and nose can occur from the day after vaccination, sometimes accompanied by wheezing, sneezing and/or coughing, particularly in very young susceptible puppies. Signs are generally transient, but in occasional cases may persist for up to four weeks. In animals, which show more severe signs, appropriate antibiotic treatment may be given.

Withdrawal Period:

None

Precautions:

Allow the diluent to reach room temperature (+15 to +25°C). Shake well after addition of diluent. The contents of the vial should be used within 1 hour after reconstitution of the product.

PSC020505_02_R1
90 x 220 mm

Shelf life: 27 months

Storage:

Vaccine and diluent should be stored between +2 °C and + 8 °C.
Keep out of reach of children.

Presentation:

1 dose, 5 dose and 10 dose vial

Manufactured by:

Intervet International B V
Wim de Körverstraat 35
5831 AN Boxmeer,
The Netherlands.

Imported and Marketed by:

Intervet India Pvt. Ltd.
Sagar complex, Bldg.B-1
Gala No. 12 to 16
Ovali Village, Bhiwandi
Thane 421302

