


Canine Leptospirosis Vaccine, Inactivated I.P.

Nobivac® LEPTO



**NOT FOR HUMAN USE; FOR ANIMAL
TREATMENT ONLY**



Warning: To be sold by retail on the prescription of a "Veterinary Doctor" only.

Composition:

Each dose of inactivated vaccine contains,
Leptospira interrogans serotype Canicola strain Ca-12-000 at least 957 units/ml and
Icterohaemorrhagiae strain 820k at least 625 units/ml

Indications:

For the active immunization against leptospirosis in dogs caused by the
Leptospira interrogans serogroup *Canicola* and *Icterohaemorrhagiae*.
Duration of immunity : 1 year

Vaccination Programmes:

Basic vaccination:

All dogs not previously vaccinated should be vaccinated twice 2-4 weeks apart.
Puppies should be at least 8 weeks of age before they receive the first vaccination.

Revaccination:

Annually

Administration:

- Shake well before use.
- Vaccines should be administered by subcutaneous injection in the neck or chest region, observing the usual aseptic precautions.

Withdrawal Period:- Not Applicable

Contra-indications and warning:

- Only healthy dogs should be vaccinated and an adequate clinical examination should be made prior to inoculation.
- Allow the vaccine to reach ambient temperature (15-25 deg. celcius) before use.
- Sterile equipment should be used for the administration, but avoid contamination of vaccine with traces of disinfectant or spirit.
- Hypersensitivity reactions following administration are rare but as with all vaccine may occasionally be encountered. In this event, administration of adrenaline injection by the subcutaneous route may be indicated.
- May be used in pregnant bitches

Compatibility:

Nobivac Lepto is compatible with:

- Nobivac DHPPi - Nobivac Parvo-C
- Nobivac DHP - Nobivac Pi
- Nobivac Puppy DP

Nobivac Lepto can also be administered with Nobivac Rabies at the same time but at a different administration site.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above.

A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Shelf life: 21 months when stored at 2°C and 8°C.

Storage:

- Store in dark between 2°C and 8°C.
- Avoid freezing.
- Care should be taken to avoid prolonged or repetitive exposure to high ambient temperatures.
- Broached vials should be used within one working day.

Manufactured by:

Intervet International BV
Boxmeer, The Netherlands.

Imported & Marketed by:

Intervet India Pvt. Ltd.
Sagar complex, Bldg.B-1
Gala No. 12 to 16
Ovali Village, Bhiwandi
Thane 421302

