

FOR THE USE ONLY OF A VETERINARIAN OR
A HOSPITAL OR A LABORATORY OR A FARM

374261 R1

Combined Canine Distemper & Canine Parvovirus Vaccine, Freeze dried including Diluent



Nobivac®
PUPPY DP



NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

WARNING: To be sold by retail on the prescription of a Registered Veterinary Practitioner only

DESCRIPTION

This vaccine contains freeze-dried living, attenuated, strains of canine distemper virus (CDV strain Onderstepoort) and canine parvovirus (CPV strain 154) grown in cell-line tissue culture. It is presented as a freeze-dried plug in single dose vials containing not less than $10^{5.0}$ TCID₅₀ CDV and $10^{7.0}$ TCID₅₀ CPV vaccine virus. Each vial contains a single dose for reconstitution. The vaccine also contains stabilizer and traces of antibiotics.

INDICATIONS

For active immunization of young puppies against canine distemper (CDV) and canine parvovirus disease (CPV).

VACCINATION PROGRAMMES

Nobivac Puppy DP is intended for use in animals with high levels of MDA against canine distemper and/or canine parvovirus. The earliest age at which these levels do not interfere with vaccination is 6 weeks in the majority of animals. To ensure protection for individual puppies that do not respond at this age due to very high levels of MDA and to provide protection against hepatitis (CAV1), CAV2 infections and parainfluenza, revaccination with Nobivac DHPPi is advised.

The following recommendations are included since it is likely that immunization with Nobivac Puppy DP will perform part of a more comprehensive vaccination programme:

a. Programme for pups where exposure to distemper and/or parvo is possible before 8-10 weeks of age; and the MDA status of the pup is unknown:

Age 4-6 weeks - Nobivac Puppy DP

Age 8-10 weeks - Nobivac DHPPi with Nobivac Lepto or Nobivac L4

Age 12 weeks - Nobivac DHPPi with Nobivac Lepto or Nobivac L4, Nobivac Rabies or Nobivac RL

b. Programme for pups where vaccination is started at the age of 8-10 weeks:

Age 8-10 weeks - Nobivac DHPPi with Nobivac Lepto or Nobivac L4

Age 12 weeks - Nobivac DHPPi with Nobivac Lepto or Nobivac L4, Nobivac Rabies or Nobivac RL

c. Programme for pups where vaccination is not started before the age of 12 weeks:

Age 12 weeks - Nobivac DHPPi with Nobivac Lepto or Nobivac L4, Nobivac Rabies or Nobivac RL

Age 14-16 weeks - Nobivac Lepto or Nobivac L4

PSC020502_02_R1

90 x 160 mm

REVACCINATION

It is recommended that dogs be revaccinated against:

1. Canine distemper, canine infectious hepatitis, canine parvovirus disease and respiratory disease caused by canine adenovirus type-2 - every 3 years.
2. Leptospirosis and respiratory disease caused by canine parainfluenza virus- every year.
3. Rabies - every 3 years (can be modified to comply with local regulations).

DOSAGE AND ADMINISTRATION

The contents of one vial of reconstituted vaccine should be injected subcutaneously. Reconstitute immediately prior to use by the addition of the contents of one vial (1.0 ml) of Nobivac Solvent. After reconstitution, the vaccine should be used within 30 minutes.

RECOMMENDATIONS FOR USE

Only healthy dogs should be vaccinated and an adequate clinical examination should be made prior to inoculation. Sterile equipment should be used for administration, but avoid contamination of vaccine with traces of disinfectant or spirit.

CONTRA-INDICATIONS AND WARNINGS

Occasionally a mild anaphylactic type hypersensitivity reaction may occur after vaccination, as is possible after foreign proteins. Such reactions are, in most cases, self limiting.

SHELF LIFE : 24 months

PHARMACEUTICAL PRECAUTIONS AND STORAGE

Store in dark between 2°C and 8°C. Care should be taken to avoid prolonged or repetitive exposure to high ambient temperatures following withdrawal from the refrigerator prior to use; under hot summer conditions vaccine potency can be severely reduced within a few hours. Nobivac Solvent may be stored at room temperature.

PRESENTATION

Boxes containing vials of Nobivac Puppy DP and vials of Nobivac Diluent, or boxes containing only vials of Nobivac Puppy DP.

PACKAGE QUANTITY

Cartons of 5, 10, 25 or 50 single dose vials.

FURTHER INFORMATION

Experience has shown that the maternal antibody status of pups within a litter varies greatly and reliance should not be placed on serological examination of the bitch alone.

Manufactured by:

Intervet International BV
Boxmeer, The Netherlands

Imported and Marketed by:

Intervet India Pvt. Ltd.
Sagar complex, Bldg. B-1
Gala No. 12 to 16
Ovali Village, Bhiwandi
Thane 421302

