

RANIKHET DISEASE VACCINE, LIVE, I.P.

Nobilis®
ND CLONE 30



Warning: To be sold by retail on the prescription of a 'Veterinary Doctor' only

NOT FOR HUMAN USE, FOR ANIMAL TREATMENT ONLY.

COMPOSITION

Nobilis ND Clone 30 is a live, freeze-dried vaccine against Newcastle Disease (ND). Each dose contains at least 10^6 ELD₅₀ of the ND strain Clone 30. The vaccine pellet contains stabilizers.

INDICATION

Immunization of chickens and turkeys against Newcastle Disease.

ADMINISTRATION

The vaccine can be administered by spray, by ocular or nasal instillation or through the drinking water. For the ocular/nasal instillation a special diluent, Diluent Oculo-Nasal is available.

VACCINATION PROGRAM

The vaccine is safe to use from 1 day of age onwards. The optimum time and method of the first administration and re-vaccination depends largely upon the local situation. Therefore, the advice of a veterinary surgeon should be sought.

The vaccine can be administered to 1-day old chicks and older chickens by coarse spray or by the intranasal/ocular route of administration. The vaccine can be administered to 7-day and older chicks by drinking water.

IMMUNITY

An adequate immunity against ND will last for approximately 6 weeks in chickens and 4 weeks in turkey. For longer protection re-vaccination is necessary.

VACCINATION REACTION

Duration and intensity of the vaccination reaction are generally dependent on the health and condition of the birds. Hygiene and management are also important in the post vaccination period. On the basis of experience with Nobilis ND Clone 30 emergency, vaccination during lay is not expected to have a negative influence on the egg production when administered to animals that have been previously vaccinated with live and inactivated Newcastle disease vaccine.

WITHDRAWAL PERIOD

None.

LIST of EXCIPIENTS:

- Sorbitol
- Hydrolysed gelatine
- Pancreatic digest of casein

- Disodium phosphate dihydrate
- Water for injections

PRECAUTIONS

- The ND virus can cause conjunctivitis in man. Wash and disinfect hands and equipment after vaccinating. Avoid exposure to vaccine virus as much as possible.
- Any surplus vaccine should be destroyed by burning or by boiling.
- Vaccinate healthy birds only.
- Each vial should be used immediately after opening.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

Safety and/or efficacy data are available which demonstrate that this vaccine can be mixed with the live infectious bronchitis (strain H120 and Ma5) vaccines by Intervet.

SHELF LIFE : 24 months

STORAGE CONDITIONS

Store in a dark place between +2° and +8°C.

PRESENTATIONS

Vials of 1000, 2500 and 5000 doses in packages of 10 vials.

Imported & marketed by:

Intervet India Pvt. Ltd.
Sagar complex, Bldg.B-1,
Gala No.12 to 16,
Ovali Village, Bhiwandi,
Thane 421302 India.

Manufactured by:

Intervet International BV
Wim de Körverstraat 35,
5831 AN
Boxmeer, The Netherlands
Imp. Lic. No.: SV-6-597

