

FOR THE USE ONLY OF A VETERINARIAN OR A HOSPITAL OR A LABORATORY OR A FARM

# Pheniramine Maleate Injection I P

**Avilin<sup>®</sup>** VET



**NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY**

SCHEDULE G PRESCRIPTION DRUG - CAUTION It is dangerous to take this preparation except under veterinarian's supervision

## COMPOSITION

Each ml contains:

Pheniramine Maleate I P ..... 22.75 mg

As one of many antihistaminic compounds, Avilin<sup>®</sup> is noteworthy for its potent, rapid and prolonged action as well as for its excellent tolerance.

## INDICATIONS

An allergic reaction is assumed to be the cause of the following diseases or symptoms in veterinary medicine:

Itching of unknown genesis and of various localisation, eczema, dermatitis, urticaria, skin oedema, insect bites, photodermatitis, rhinitis, tail eczema in horses, stomatitis, toxic hoof corns and inflammation of the hooves of cattle, serum shock, paresis during pregnancy, puerperal toxemia and secondary retention, pulmonary oedema in cattle, pulmonary emphysema in horses, toxic hypopepsia.

The efficacy of Avilin<sup>®</sup> has also been demonstrated in drug

allergies, anaphylactic shock, laminitis, bloat in ruminants due to histamine atony of the rumen and also in the treatment of acute septic metritis in cows in conjunction with anti-bacterial therapy.

## DOSAGE AND ADMINISTRATION

The dosage should be adjusted individually. If possible, small doses are given first because it has often been shown that antiallergic drugs are effective in comparatively smaller dosage.

Large animals : 5 - 10 ml or more

Small animals : 1/2 - 1 ml or more

Avilin<sup>®</sup> may be administered intramuscularly or intravenously. The onset of action depends on the mode of administration. Mostly there is an immediate effect on intravenous administration. On intramuscular administration, it takes about 10-20 minutes for the effect to set in.

Injection may be repeated after 8-12 hours, if necessary.

**STORAGE :** Store protected from light

**PRESENTATION :** Vial of 10 ml, 33 ml

**WITHDRAWAL PERIOD :** Meat - 7 days, Milk - 2 days

® Registered Trademark

Marketed by :

**Intervet India Pvt. Ltd.**

Sagar Complex, Bldg., B-1 Gala No. 12-14,  
Ovali Village, Bhiwandi, Thane - 421 302

Manufactured by : Samrudh Pharmaceuticals Pvt. Ltd.  
J-174, J-168, J-168/1, MIDC, Tarapur, Thane - 401506

 **MSD**  
Animal Health