

FOR THE USE ONLY OF A VETERINARIAN OR A HOSPITAL OR A LABORATORY OR A FARM



Prednisolone Acetate Injectable Suspension U.S.P.



NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG CAUTION Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

COMPOSITION

Each ml contains:

Prednisolone Acetate I.P. 10 mg

Benzyl Alcohol I.P. 0.945 % v/v

PREDNISOLONE Crystalline injectable suspension was developed for use in the veterinary field. It has proved to be highly effective in the treatment of acetonemia in cattle (bovine ketosis). Following the injection of PREDNISOLONE Crystalline Suspension, the blood glucose level is restored to normal in short time, appetite and milk yield improves, and circulating ketone bodies return to a normal level. It is three to five times more active than hydrocortisone, (natural cortizones) but does not affect mineral balance as that of the other potent corticosteroids.

INDICATIONS

PREDNISOLONE Crystalline Suspension is used in treating ketosis in dairy cattle, rheumatoid arthritis, bursitis inflammatory and allergic conditions in large and small animals in conjunction with appropriate antibiotic or chemotherapeutic agent when indicated.

DOSE AND ADMINISTRATION

For Systemic treatment, injection is given deep into the muscle, 2-4 times if necessary at an interval of 24 hours.

Cattle : 50 - 100 - 200 mg

Cows : 25 - 50 mg

Piglets, dogs, cats : 10 - 30 mg

For local treatment injection is given in the part involved carefully under strict aseptic conditions. Contra indications include diabetes mellitus & osteoporosis. In such cases, Crystalline suspension must not be administered more than twice.

Injection into Joints & tendon sheaths

Large animals : 25 - 75 mg or more

Dogs & Cats : 5 - 20 mg or more

Injection into Bursae

10 - 50 mg or more

Subconjunctival injection

1.25 - 12.5 mg

Retrobular injection

2.5 - 12.5 mg

Infiltration (as in periostitis etc.)

10 - 50 mg

Injection into infected tissues are contra-indicated. The preparation should be shaken well before use. Use fine needle for local therapy.

STORAGE : Store protected from light

WITHDRAWAL PERIOD :

Meat - Cattle, Sheep & Goat - 5 days & Pig - 28 days

Milk - 3 days

PRESENTATION :

Vial of 10 ml

Made in India by :

Intervet India Pvt. Ltd.,
Sagar Complex, Bldg., B-1 Gala No. 12-14,
Ovali Village, Bhiwandi, Thane - 421 302

Manufactured at : Gat No. 285 - B, Kasaramboli,
Pune - 411 004

