

FOR THE USE ONLY OF A VETERINARIAN OR A HOSPITAL OR A LABORATORY OR A FARM

Rx  
**Levamisole Hydrochloride & Oxytoclanide Suspension**  
**Tolzan® Plus-L VET**

**Fluke & Worm Drench**



**NOT FOR HUMAN USE-FOR ANIMAL TREATMENT ONLY**

**SCHEDULE H PRESCRIPTION DRUG - CAUTION** Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

**COMPOSITION :**

Levamisole Hydrochloride IP ..... 2.5% w/v  
Oxytoclanide IP ..... 3.4% w/v

**INDICATIONS:**

For the treatment and control of round worms and liver flukes in cattle, sheep and goats.

**Liver fluke**

Controls both adult liver fluke (*Fasciola hepatica*) and adult giant liver fluke (*Fasciola gigantica*) in cattle, sheep and goats.

**Conical fluke**

Controls immature and adult conical fluke (*Paramphistomum spp.*) in cattle, sheep and goats.

**Targets :** Wireworm, Brown Stomachworm, Bankruptworm, Cattle Bankruptworm, Long-necked Bankruptworm, Large-Mouthed Bowelworm, Hookworm, Nodularworm, Lungworm

**PRECAUTIONS & WARNINGS:**

- No worm remedy, not even Tolzan Plus-L, should be dosed to sheep and goats that have not been immunized against Pulpy Kidney (Enterotoxaemia).
- In a small percentage of high-producing dairy cows the following transient reactions may be noted: softening of the faeces, salivation and swelling of the head.

- In case of poisoning call a doctor and make this label available to him.
- KEEP OUT OF REACH OF CHILDREN AND UN-INFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice.

**SAFETY:**

Tolzan Plus-L may be dosed to all cattle, sheep and goats without any pre-dosing, starving or change of diet.

**DIRECTIONS FOR USE:**

- Use as directed
- Shake well before use.

**DOSAGE:**

Cattle : 30 ml / 100 kg live body weight  
Sheep and goats : 3 ml / 10 kg live body weight  
For the treatment of conical fluke (*Paramphistomum spp.*) : 3 ml / 10 kg live body weight, followed by a second identical dose 72 hours later.

**ROUTE OF ADMINISTRATION:**

Oral

**STORAGE:**

Store protected from light

**PRESENTATION:**

120 ml plastic bottle, 1 Ltr jerry can

**WITHDRAWAL PERIOD :**

Meat - 14 days  
Milk - 07 days

Marketed by :

**Intervet India Pvt. Ltd.**  
Sagar Complex, Bldg., B-1 Gole No. 12-14,  
Ovali Village, Bhiwandi, Thane - 421 302

Manufactured by : Medibios Laboratories Ltd.,  
Plot No. J76, M.I.D.C., Tarapur, Thane - 401 506  
at - N - 50, M.I.D.C., Tarapur, Thane - 401 506

