

FOR THE USE ONLY OF A VETERINARIAN OR A HOSPITAL OR A LABORATORY OR A FARM

℞ **Injection of Analgin**
Vetalgin® VET



Analgesic, Antispasmodic, Antirheumatic and Antipyretic

NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG - CAUTION : Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

COMPOSITION

Each ml contains :

Analgin IP 0.5 g
Chlorbutol IP (as bacteriostatic) 0.4 % w/v

CHEMISTRY AND PHARMACOLOGY

VETALGIN® belongs to the series of the phenyldimethyl-pyrazolone derivatives.

VETALGIN® has a central analgesic action and at the same time marked sedative action on animals.

The spasmolytic effect on smooth muscle is combined with peripheral vasodilation, producing an increased blood supply to striped muscle which is therapeutically desirable,

VETALGIN® causes no adverse effects on circulation, heart, kidneys or other organs at therapeutic dose. It is likewise well tolerated by young animals.

INDICATIONS

IN ALL ANIMALS :

For relief of pain in colic and other spastic conditions of the abdominal organs; acute and chronic arthritis, rheumatic conditions of muscles and joints, neuritis, neuralgia, tendo-vaginitis. Obstruction of the oesophagus by foreign bodies - to relieve spasm. In gastritis of small domestic animals to prevent vomiting.

IN THE HORSE :

For relief of pain, relaxation of intestinal spasm, for careful stimulation of peristalsis in colic, in acute stomach dilation, to produce calmness during examination or treatment, in tetanus for relieving spasm (combined with tetanus antiserum).

IN THE PIG : Spasm of the cervical orifice during parturition.

Serositis and inflammation of the joints in piglings.

CONTRAINDICATIONS

- Do not use in cats.
- Do not use by subcutaneous route.
- VETALGIN® should not be used in animals with disturbances of haematopoietic system

ADMINISTRATION

By intravenous injection in acute condition or by deep intramuscular injection. Combination of both kinds of injections is possible. Injections can be repeated without hesitation on the same day.

Subcutaneous injection is not recommended as it may cause irritation of the tissue

DOSE

Cattle : 8 ml per 100 kg body weight
Horses : 20-60 ml
Calves : 5-15 ml
Sheep and Goats : 2-8 ml
Pigs : 10-30 ml (Spasm of cervical orifice : 5-10 ml i.m.)
Dog : 1-5ml

STORAGE : Store protected from light

PRESENTATION : Vial of 30 ml, 33 ml

WITHDRAWAL PERIOD : Meat - Cattle-12 days, Pig-3 days, Sheep & Goat-18 days
Milk - Cattle & Pig 2 days or 4 milkings, Sheep & Goat 3 days or 6 Milkings

Marketed by : **Intervet India Pvt. Ltd.**

Briahnagar, Off Pune-Nagar Road,
Wagholi, Pune - 412 207, India

Manufactured by : Samrudh Pharmaceuticals Pvt. Ltd.
J-174, J-168, J-168/1, MIDC, Tarapur, Thane - 401506

