



347857 R1



INTRAMAMMARY SUSPENSION IN PREFILLED SYRINGE

YELINTRA * TM**VETERINARY****Composition:**

Per syringe of 8gm contains:

Tetracycline HCL	200 mg
Neomycin base (as sulphate).....	250 mg
Bacitracin	2000 IU
Prednisolone	10 mg

NOT FOR HUMAN USE; FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG – CAUTION Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

Target Species:

Lactating dairy cattle.

Indications:

Intramammary administration to lactating cows for therapeutic treatment of clinical mastitis caused by sensitive microorganisms (*S. aureus*, *Streptococci*, *E.coli*, *Klebsiella spp.*, *A. pyogenes*)

Posology and method of administration:

Intramammary administration of one syringe per infected quarter every 12 hours. Maximum treatment is four administrations.

Before use, milk out the quarter completely. Partially or totally, remove the cap from the end of the nozzle of the injector, depending on partial or total insertion. Insert the nozzle carefully into the teat opening. Squeeze the entire contents of the injector slowly into the teat.

Contraindications:

Do not use in animals known to be allergic to tetracycline, bacitracin and neomycin.

Do not use the cleaning towels on teats with visible not healed cuts or wounds.

Adverse Reaction:

None observed. If you notice any side effects, please contact your veterinary surgeon.

Special precautions for use in animals:

- The product is intended to be used during lactation and can be used during pregnancy.
- Inappropriate use of the product may increase the prevalence of bacteria resistant to tetracycline, bacitracin or neomycin.

Special precautions to be taken by the person administering the medicinal product to animals:

- Individuals allergic to tetracyclines, neomycin or bacitracin should avoid skin contact with the product.
- Tetracycline, neomycin and bacitracin may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Allergic reactions to these substances may occasionally be serious.
- If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the Doctor this warning. Swellings of the face, lips or eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Use during pregnancy or lactation:

The product is intended to be used during lactation and pregnancy. Studies in laboratory animals have not produced any evidence of embryotoxic and/or teratogenic effects.

Withdrawal Period(s):

Meat and offal: 14 days
Milk: 96 hours (8 milkings)

Shelf Life: 24 months**Special Storage Conditions:**

Store in a refrigerator (2°C to 8°C)

Package Size:

Box of 1 sachet of 4 syringes and 4 cleaning towels
Box of 1 sachet of 20 syringes and 20 cleaning towels

Manufactured by:

Intervet International B.V.
Wim de Korverstraat 35,
5831 AN Boxmeer,
The Netherlands

Imported & Marketed by:

Intervet India Pvt. Ltd.
Sagar complex, Bldg.B-1
Gala No. 12 to 16
Ovali Village, Bhiwandi
Thane 421302

*TM- Under Registration



