Infectious Chicken Anemia Vaccine, Live

232240 R1

Nobilis®
CAVP4

NOT FOR HUMAN USE FOR ANIMAL TREATMENT ONLY

WARNING: To be sold by retail on the prescription of a "Veterinary Doctor" only.

Description
Nobilis® CAVP4 is a Lyophilisate and solvent for suspension for injection for chicken against chicken anemia virus

Composition
Each dose contains:

Active Ingredients	**Quantity**
Composition of vaccine before freeze drying	
CAA strain 26P4 infected embryo homogenate	0.67 ml
Composition of vaccine after freeze drying	
Live Attenuated CAA virus strain 26P4	At least 3.0 $\log_{10}$ $TCID_{50}$ per bird dose
Inactive Component	
Pancreatic digest of casein	6.7 mg
Dextran 70	13.3 mg
Sorbitol	13.3 mg
Sucrose	25 mg
Gelatin	8.3 mg
Dibasic Potassium Phosphate	0.4 mg
Monobasic potassium Phosphate	0.2mg
Gentamicin sulphate	0.1 mg
Water for Injection	1.0ml

Solvents:
Dilavia (for intramuscular/subcutaneous injection)
Adjuvant:
dl-a-tocopherol: 75 mg/ml

Unisolve (for intracutaneous application via wingweb)

Indication
Used for passive immunization of chicks, via active immunization of breeders, against afflictions caused by chicken anemia virus.

Target Species
Chicken

Contraindications
None

Special warnings
None.

Special precautions for use
Special precautions for use in animals
Vaccinate only healthy animals

Special precautions to be taken by the person administering the veterinary medicinal product to animals
In the case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Adverse reactions (frequency and seriousness)
None

Use during pregnancy, lactation or lay
Not to be used during the laying period.

Interaction with other medicinal products and other forms of interaction
No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other medicinal product therefore needs to be made on a case-by-case basis.

Moved to repository by TORALY 13:Mar:2025 13:20

Method of administration
One dose per bird via intramuscular, subcutaneous or intracutaneous via Wing-web administration.

Intramuscular or subcutaneous administration:
Reconstitute the vaccine in the diluent Dilavia. Select the amount of diluent to be used on the number of doses per vial (200ml per 1000 doses). Administer 0.2 ml of the vaccine by intramuscular or subcutaneous injection.

Intracutaneous administration via wing-web method:
Reconstitute the vaccine in the diluent Unisolve. Select the amount of diluent to be used on the number of doses per vial (13ml per 1000 doses) Dip the double needle before vaccination of each bird into the vaccine suspension so that both grooves are filled. Insert the needle from beneath through the wing-web.

Vaccination scheme:
Single vaccination with one dose per bird from the age of 6 weeks, but at least 6 weeks before the start of the lying period.

Overdose (symptoms, emergency procedures, antidotes), if necessary
No symptoms at a ten-fold overdose.

Withdrawal period(s)
Zero days

Pharmacotherapeutic group: Poultry, live viral vaccine.
The P4 strain is an attenuated strain of the Chicken Anemia Virus, with good immunogenic properties but a significantly reduced pathogenicity for day-old chickens. Vaccination of the breeder flock before the onset of lay induces high uniform levels of neutralizing antibodies, which prevent vertical transmission of virulent CA V to the progeny. In the progeny, the high levels of MDA prevent clinical disease during the first critical weeks of their lives.

List of excipients
Pancreatic digest of casein
Dextran 70
Sorbitol
Sucrose
Gelatin
Dibasic Potassium Phosphate
Monobasic Potassium Phosphate
Gentamicin Sulphate
Water for injections

Incompatibilities
Do not mix with any other veterinary medicinal product, except the diluent supplied for use with the product.

Shelf life
Shelf-life of the lyophilisate as packaged for sale: 24 months
Once reconstituted use within 4 hours

Pack size and Presentation
- **Freeze dried pellet**: Glass vials of 1000 doses.
- **Diluent Dilavia**: 200 ml in glass or PET vial
- **Diluent Unisolve**: 13ml in glass vial

Special Storage precautions
Freeze dried pellet: Store at +2°C to +8°C. Do not freeze. Protect from light.
Solvent: Store below 25°C, if stored independently from the vaccine.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products
Disinfect and wash hands and vaccination equipment after vaccination.
Dispose of waste material by boiling, incineration or immersion in an appropriate disinfectant, in accordance with the national requirements.

Manufactured by:
Intervet International B.V.,
Wim de Korverstraat 35, 5831
AN Boxmeer, The Netherlands.

Imported & Marketed by:
Intervet India Pvt. Ltd., Bldg. No. B-1, Godown No.12 to 16, Ground Floor & Godown No.12-A to 17-A First Floor, House No. 233, Sagar Complex, Owali, Bhiwandi, Thane -421302,Maharashtra, India.

MSD
Animal Health

PSC020505 02 R1 90 x 220 mm Black

Moved to repository by TORALY 13:Mar:2025 13:20