

FOR THE USE ONLY OF A VETERINARIAN OR
A HOSPITAL OR A LABORATORY OR A FARM

INFECTIOUS CORYZA VACCINE, INACTIVATED IP
Nobilis®
CORYZA

NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY-



Warning: To be sold by retail on the prescription of a 'Veterinary Doctor' only

COMPOSITION

Each 0.25 ml dose contains:

Active Ingredients	Quantity
A. <i>paragallinarum</i> - Strain 083(Serotype A) - Strain Spross (Serotype B) - Strain H-18 (Serotype C)	At least 1 CPD ₇₀ At least 1 CPD ₇₀ At least 1 CPD ₇₀
Excipient	
Purified saponin	500 µg
KH ₂ PO ₄	0.1 mg
Na ₂ HPO ₄	0.3 mg
NaCl	1.7 mg
AF 9030E	0.04 mg
Water for injections	250 mg

INDICATION

Vaccination against *Avibacterium paragallinarum* infections in chickens.

VACCINATION PROGRAM

Chickens can be vaccinated from 10 weeks of age onwards. To achieve a protection in layers and breeders of 52 weeks post vaccination,two vaccinations are required. The first dose has to be administered at the age of 10-12 weeks and the repeated dose a few weeks before the onset of egg production. The interval between two vaccinations has to be at least 4 weeks.

DOSAGE AND ADMINISTRATION

Intramuscular injection of 0.25 ml of vaccine in chickens from 10 till 12 weeks old and a revaccination ultimately several weeks before lay. Interval between vaccinations at least 4 weeks.

VACCINATION REACTION

In healthy animals no abnormal clinical reactions are observed. Slight transient swelling at the site of injection for some weeks after vaccination is occasionally observed.

CONTRAINDICATIONS

None

USE DURING LAY

Use during lay is not recommended.

INTERACTIONS WITH OTHER MEDICAMENTS AND FORMS OF INTERACTION

Do not administer in conjunction with other systemic treatments.

OVERDOSE

No particular symptoms at double dose

WITHDRAWAL PERIOD

Zero days.

NOTE

- Vaccinate healthy birds only.
- Before using vaccine allow it to reach room temperature (15 °-25 °C).
- Shake bottle prior to and intermittently during application.
- Use sterilized syringe and needle.
- Opened bottles should be used within 3 hours.
- Should a vaccinator by accident inject himself or a bystander, a local reaction may occur. It is recommended that the advice of a doctor be sought, taking care to inform the doctor about the nature of vaccine.

SHELF LIFE : 24 months for Polyethylene terephthalate (PET) & 12 months for glass bottles when stored at 2 to 8 °C

STORAGE CONDITIONS

Store in dark between 2 °C and 8 °C.
Avoid freezing.

Manufactured by:

Intervet International B.V.
Wim de Korverstraat 35
5831 AN Boxmeer, The Netherlands.

Imported and Marketed by:

Intervet India Pvt. Ltd., Bldg. No. B-1, Godown No.12 To 16, Ground Floor & Godown No. 12-A To 17-A First Floor, House No. 233, Sagar Complex
Owaji, Bhiwandi, Thane 421302, Maharashtra, India.



