

RANIKHET DISEASE VACCINE INACTIVATED IP

Nobilis®

ND BROILER



NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

Warning: To be sold by retail on the prescription of a 'Veterinary Doctor' only

DESCRIPTION

Inactivated vaccine for the protection of day old chickens against Newcastle Disease. Nobilis ND Broiler contains one immunogenic strain of Newcastle Disease virus. The virus has been grown on embryonated eggs and is inactivated with formalin. Subsequently they have been suspended in the aqueous phase of an oil adjuvant emulsion.

COMPOSITION

Composition:

Each 0.1 ml dose contains:

Active Ingredients	Quantity
Inactivated Newcastle Disease Virus* (strain Clone 30)	Containing ≥ 20 PD ₅₀ units/dose (protective dose 50%) or inducing ≥ 4.0 log ₂ HI units per 1/50 th dose
Excipients	
Light liquid paraffin	552 mg/ml
Polysorbate 80	6.2 mg/ml
Sorbitan mono-oleate	51 mg/ml
Glycine	4.8 mg/ml
Formaldehyde	<0.05%
Water For Injection	q.s. to 1.0 ml
* According to potency test	

INDICATION

For the vaccination of chickens against disease caused by Newcastle disease virus.

VACCINATION PROGRAMME

Nobilis ND Broiler is administered by injection at day-old, in combination with live ND vaccine, e.g. Nobilis ND Clone 30, administered by spray or the oculo-nasal route.

DOSAGE AND ADMINISTRATION

Inject 0.1 ml per broiler intramuscularly into the thigh muscle, or subcutaneously into the lower part of the neck.

VACCINATION REACTION

In healthy broilers no clinical reaction to vaccination will be observed. For some weeks after vaccination a very slight swelling may occur at the site of injection. Local tissue reaction may occur.

IMMUNITY

Healthy broilers, vaccinated at day-old with Nobilis ND Broiler and Nobilis ND Clone 30 will be protected until slaughter.

CONTRA-INDICATIONS

When used according to instructions: None

UNDESIRABLE EFFECTS

In healthy animals no clinical reactions. Slight swelling at the site of injection for some weeks after vaccination is occasionally observed.

Use during lay: Not Applicable

NOTE

- Only vaccinate healthy broilers.
- Before using vaccine allow it to reach room temperature (15-25 °C).
- Shake the bottle before use.
- Use sterilised syringe and needle.
- Use the entire contents when first opened.
- Do not mix with other vaccines.

SPECIAL PRECAUTIONS FOR USE:

Special precautions to be taken by the person administering the veterinary medicinal product to animals

To the user:

This veterinary medicinal product contains mineral oil. Accidental injection/self-injection may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If you are accidentally injected with this product, seek prompt medical advice even if only a very small amount is injected and take the package leaflet with you. If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This veterinary medicinal product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

WITHDRAWAL PERIOD: NONE

SHELF LIFE

36 months

STORAGE: Store in dark between 2 °C and 8 °C.

PACK: Bottles containing 200 ml sufficient for 2000 doses.

Manufactured by:

Intervet International B.V.
Wim de Korverstraat 35
5831 AN Boxtmeer
The Netherlands.

Imported & Marketed by:

Intervet India Pvt. Ltd., Bldg. No. B-1, Godown No.12 to 16, Ground Floor & Godown No.12-A to 17-A First Floor, House No. 233, Sagar Complex, Owali, Bhiwandi, Thane -421302, Maharashtra, India.



