

235566 R1

**Canine Leptospirosis Vaccine,
Inactivated I.P.
Nobivac® LEPTO**



**NOT FOR HUMAN USE; FOR ANIMAL
TREATMENT ONLY**



Warning: To be sold by retail on the prescription of a "Veterinary Doctor" only.

Composition:

Each dose of inactivated vaccine contains,
Leptospira interrogans serotype Canicola strain Ca-12-000 ≥ 957 Units/ml
Leptospira interrogans serotype Icterohaemorrhagiae strain 820K ≥ 625 Units/ml

Modified Hartmann's solution

NaCl.....6mg

KCl.....0.4mg

L-lactic acid sodium salt..... 4.6mg

CaCl₂·2H₂O.....0.36mg

Water for Injections.....1 ml

Indications:

For the active immunization against leptospirosis in dogs caused by the
Leptospira interrogans serogroup Canicola and Icterohaemorrhagiae.

Amounts to be administered and administration route:

Shake well before use.

Vaccines should be administered by subcutaneous injection in the neck or chest region,
observing the usual aseptic precautions.

Subcutaneous injection of 1 ml per animal.

Basic vaccination:

All dogs not previously vaccinated should be vaccinated twice 2-4 weeks apart. Puppies
should be at least 8 weeks of age before they receive the first vaccination.

Revaccination:

annually.

Duration of immunity: 1 year

Contraindications:

None

Special warnings:

None

Special precautions for use:

Allow the vaccine to reach room temperature (15-25°C) before use.

Sterile injection equipment should be used.

Special precautions for use in animals:

Vaccinate healthy animals only.

**Special precautions to be taken by the person administering the veterinary
medicinal product to animals:**

None

Adverse reactions (frequency and seriousness):

An occasional, mild, self-limiting hypersensitivity reaction may occur.

A local reaction of limited size may arise during the first few days after vaccination. Such reaction may evolve to a more severe condition (anaphylaxis), which may be life-threatening. In very rare cases, clinical signs of immune-mediated haemolytic anaemia, immunemediated thrombocytopenia, or immune-mediated polyarthritis have been reported.

Use during pregnancy and lactation:

Can be used in pregnant bitches.

Overdose (symptoms, emergency procedures, antidotes) (if necessary):

No particular symptoms at double dose.

Note that product is presented in single dose vials.

Withdrawal period(s) : Not applicable.

Incompatibilities

Do not mix with any other veterinary medicinal product except the vaccines of the Nobivac series mentioned under compatibility.

Compatibility:

Nobivac Lepto is compatible with:

- Nobivac DHPPi - Nobivac Parvo-C
- Nobivac DHP - Nobivac Pi
- Nobivac Puppy DP

Nobivac Lepto can also be administered with Nobivac Rabies at the same time but at a different administration site.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

List of excipients:

- NaCl
- KCl
- L-Lactic acid sodium salt
- $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$
- Water for injections

Shelf life: 21 months when stored at 2°C and 8°C.

Storage:

- Store in dark between 2°C and 8°C.
- Avoid freezing.
- Care should be taken to avoid prolonged or repetitive exposure to high ambient temperatures.
- Broached vials should be used immediately.

Manufactured by:

Intervet International B.V.
Wim de Korverstraat 35
5831 AN Boxmeer
The Netherlands.

Imported & Marketed by:

Intervet India Pvt. Ltd.
Bldg. No. B-1, Godown No.12 to 16, Ground Floor & Godown
No.12-A to 17-A First Floor, House No. 233,
Sagar Complex, Owali, Bhiwandi,
Thane -421302, Maharashtra, India.

