



NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG - CAUTION Not to be sold by retail without the prescription of a Registered Veterinary

COMPOSITION

Panacur® Powder 25% W/W · Fach gram contains DOSE FOR HORSES Fenbendazole IP 250 mg Excipients Panacur® 150 mg Tablet : EachTablet contains Fenbendazole IP 150 mg Excipients q.s. Panacur® 1.5 g Bolus : Each bolus contains body weight. 1.5 gm Fenbendazole IP Excipients Panacur® SUSPENSION (Shake well before use) Panacur® 3.0 g Bolus : Fach bolus contains Panacur® Suspension 2.5 % (90 ml, 450ml & 1 lit) Fenbendazole IP 3.0 gm Excipients q.s. Colour - Tartrazine a sheep or goat weighing 30 kg. : Each 10 ml contains Panacur® Susp. 2.5% 250 mg Fenbendazole IP

q.s.

a.s.

1.0 gm

INDICATIONS

Panacur® Susp. 10%

Broad spectrum anthelmintic against immature and mature stages of gastrointestinal nematodes, lungworms and tapeworms such as : Haemonchus spp.,Ostertagia spp.,Trichostrongylus spp.,Cooperia spp.,Nematodirus spp., Neoascaris vitulorum, Chabertia spp., Bunostomum spp., Gaigeria pachyscelis, Oesophagostomum spp., Capillaria spp., Trichuris spp., Strongyloides spp., Dictyocaulus filaria, Dictycaulus viviparus, Moniezia spp., in cattle, buffalo, sheep and goats. Infestation of dogs with Ancylostoma spp., Infestation of horses with Strongyles, Ascarids (Parascaris), Oxyuris and Strongyloides. Infestation of pigs with Hystrogylusrubidus. Oesophagostromum spp., Ascaris suum, Trichris suis and Metastronaylus spp.

Excinients

: Each 10 ml contains

Fenbendazole IP Excipients

DOSE AND ADMINISTRATION

Panacur® POWDER 25% W/W

For preparing the ready-to-use suspension, mix the contents of the pack, with slight stirring as follows:

| c pack, | will slight silling as lollows. |
|---------|---------------------------------|
| 6 g | powder with 100 ml water |
| 60 g | powder with 1 litre water |
| 120 a | nowder with 2 litre water |

Dose recommended for cattle, buffaloes, sheep, goats and pigs, Infestation with gastrointestinal nematodes and lungworms: 1 ml suspension per 3 kg body weight (i.e. 5 mg Fenbendazole per kg body weight)

| Body weight kg | Suspension ml | Body weight kg | Suspension ml |
|-------------------|------------------|-------------------|------------------|
| 15 | 5 | 120 | 40 |
| 30 | 10 | 150 | 50 |
| 45 | 15 | 210 | 70 |
| 60 | 20 | 300 | 100 |
| 90 | 30 | 450 | 150 |

The suspension is administered by means of a bottle or a drenching gun.

Panacur® TABLET

Panacur® tablet is to be administered orally by an applicator or hand. One Panacur® tablet 150mg should be administered to sheep, goats and calves weighing upto 30 kg. For every additional 30 kg body weight, one extra 150 mg Panacur® tablet to be administered.

For dogs Panacur® tablet to be administered 50 mg per kg and this is to be given for three consecutive days.

Panacur® BOLUS

Panacur® bolus 1.5 g and 3.0 g is to be given to animals at the rate of 5 mg to 10 mg per kg, body weight or as directed by veterinarian. (For treatment of heavy infestation with tapeworms, use 10 mg per kg body weight).

In horses, it is advisable to use 7.5 mg Fenbendazole per kg body weight, one 150 mg tablet should be used for each 20 kg body weight Or 1.5 g bolus for each 200 kg

6 ml of Panacur® 2.5 % Suspension is to be administered to

90 ml of $Panacur^{\otimes}$ 2.5% Suspension to be administered to a cow or a buffalo weighing 450 kg. Or as directed by a veterinarian.

20 ml of Panacur® 2.5% Suspension is to be administered to a dog weighing 10 kg and this is to be given on 3 consecutive days.

Panacur® Suspension 10 % (450 ml)

15 ml of Panacur® Suspension 10 % is to be administered to animal weighing 300 kg

PLEASE NOTE

Dietetic measures before or after treatment are not required. If reinfestation occurs, treatment should be repeated

Panacur® is well tolerated.

DDESENITATION

Panacur® POWDER

Box of 20 sachets of 6 g, 60 g and 120 g container

Panacur® TABLET / BOLUS

150 mg tablet: Box of 20 strips of 10 tablets each 1.5 g bolus: Box of 30 (15 strips of 2 boli each) 3.0 g bolus : Box of 15 boli

Panacur® SUSPENSION

2.5 % - 450 ml & 1 litre container

10 % - 450 ml container

WITHDRAWAL PERIOD:

Meat: 8 days (Cattle, Buffalo, Horses) & 14 days (Sheep, Goat)

STORAGE: Store protected from light

® Registered Trademark

Marketed by: Intervet India Pvt. Ltd. Bidg. B-1, Godown No. 12 to 16, and 12-A to 17-A, Sagar Complex, Owali Village, Bhiwandi, Thane. 421302, Maharashtra, India. Consumer Care Number: +91-20-66294700/01 E-mail: infain-india@msd.com Manufactured by: EMIL Pharmaceutical Ind. Pvt. Ltd.
Plat No. J-76 and N-50
MIDC, Tarapur, Boisar- 401506,
Dist.- Palghar, Maharashtra, India 55×190

