



NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG - CAUTION Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

COMPOSITION

Panacur® Powder 25% W/W	: Each gram contains Fenbendazole IP Excipients	250 mg q.s.
Panacur® 150 mg Tablet	: Each Tablet contains Fenbendazole IP Excipients	150 mg q.s.
Panacur® 1.5 g Bolus	: Each bolus contains Fenbendazole IP Excipients	1.5 gm q.s.
Panacur® 3.0 g Bolus	: Each bolus contains Fenbendazole IP Excipients Colour - Tartrazine	3.0 gm q.s.
Panacur® Susp. 2.5%	: Each 10 ml contains Fenbendazole IP Excipients	250 mg q.s.
Panacur® Susp. 10%	: Each 10 ml contains Fenbendazole IP Excipients	1.0 gm q.s.

Broad-spectrum anthelmintic against immature and mature stages of gastrointestinal nematodes and lungworms in cattle, buffaloes, sheep, goats, horses, pigs, dogs, and also against tapeworms in sheep and goats.

INDICATIONS

Infestation of cattle, buffaloes, sheep and goats with gastrointestinal nematodes, lungworms and tapeworms, such as: *Haemonchus spp.*, *Ostertagia spp.*, *Trichostrongylus spp.*, *Cooperia spp.*, *Nematodirus spp.*, *Neosascaris vitulorum*, *Oesophagostomum spp.*, *Chabertia spp.*, *Bunostomum spp.*, *Gaigeria pachyscelis*, *Capillaria*, *Trichuris spp.*, *Strongyloides spp.*, *Dicrocoelium filaria*, *Dicrocoelium viviparus*, *Moniezia spp.*, Infestation of dogs with *Ancylostoma spp.*, Infestation of horses with *Strongyles*, *Ascarids (Parascaris)*, *Oxyuris* and *Strongyloides* Infestation of pigs with *Hyostomylus rubidus*, *Oesophagostomum spp.*, *Ascaris suum*, *Trichuris suis* and *Metastrongylus spp.*

DOSE AND ADMINISTRATION

Panacur® POWDER 25% W/W

For preparing the ready-to-use suspension, mix the contents of the pack, with slight stirring as follows:

6 g	powder with 100 ml water
60 g	powder with 1 litre water
120 g	powder with 2 litre water

Dose recommended for cattle, buffaloes, sheep, goats and pigs, Infestation with gastrointestinal nematodes and lungworms : 1 ml suspension per 3 kg body weight (i.e. 5 mg Fenbendazole per kg body weight)

Body weight kg	Suspension ml	Body weight kg	Suspension ml
15	5	120	40
35	10	150	50
45	15	210	70
60	20	300	100
90	30	450	150

The suspension is administered by means of a bottle or a drenching gun.

Panacur® TABLET

Panacur® tablet is to be administered orally by an applicator or hand. One Panacur® tablet 150mg should be administered to sheep, goats and calves weighing upto 30 kg. For every additional 30 kg body weight, one extra 150 mg Panacur® tablet to be administered.

For dogs Panacur® tablet to be administered 50 mg per kg and this is to be given for three consecutive days.

Panacur® BOLUS

Panacur® bolus 1.5 g and 3.0 g is to be given to animals at the rate of 5 mg to 10 mg per kg. body weight or as directed by veterinarian. (For treatment of heavy infestation with tapeworms, use 10 mg per kg body weight).

DOSE FOR HORSES

In horses, it is advisable to use 7.5 mg Fenbendazole per kg body weight, one 150 mg tablet should be used for each 20 kg body weight Or 1.5 g bolus for each 200 kg body weight.

Panacur® SUSPENSION (Shake well before use)

Panacur® Suspension 2.5 % (90 ml, 450ml & 1 lit)

6 ml of Panacur® 2.5 % Suspension is to be administered to a sheep or goat weighing 30 kg.

90 ml of Panacur® 2.5% Suspension to be administered to a cow or a buffalo weighing 450 kg. Or as directed by a veterinarian.

20 ml of Panacur® 2.5% Suspension is to be administered to a dog weighing 10 kg and this is to be given on 3 consecutive days.

Panacur® Suspension 10 % (450 ml)

15 ml of Panacur® Suspension 10 % is to be administered to animal weighing 300 kg

PLEASE NOTE

Dietetic measures before or after treatment are not required. If reinfestation occurs, treatment should be repeated.

Panacur® is well tolerated.

PRESENTATION

Panacur® POWDER

6 g sachet, 60 g and 120 g container

Panacur® TABLET / BOLUS

150 mg tablet : Box of 50 (5 strips of 10 tablets each)

1.5 g bolus : Box of 30 (15 strips of 2 boli each)

3.0 g bolus : Box of 10 boli

Panacur® SUSPENSION

2.5 % - 450 ml & 1 litre container

10 % - 450 ml container

WITHDRAWAL PERIOD :

Milk : 4 days

Meat: 8 days (Cattle, Buffalo, Horses) & 14 days (Sheep, Goat)

STORAGE : Store protected from light

® Registered Trademark

Marketed by: **Intervet India Pvt. Ltd.**
Bldg. B-1, Godown No. 12 to 16, and 12-A to 17-A,
Sagar Complex, Chellai Village, Bhivandi,
Thane - 421302, Maharashtra, India.
Consumer Care Number: +91-20-66294700/01
E-mail: intervet-india@merck.com
Website: www.merck-animal-health.co.in
Manufactured by: **EMIL Pharmaceutical Ind. Pvt. Ltd.**
Plot No. J-76 and N-50
MDC, Tempco, Bolar - 401506,
Dist. - Palghar, Maharashtra, India



PH40100123 35x190 mm