

 Prednisolone Acetate
Injectable Suspension U.S.P.



NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG - CAUTION Not to be sold by retail without the prescription of a Registered Veterinary Practitioner.

COMPOSITION

Each ml contains:

Prednisolone Acetate IP	10 mg
Benzyl Alcohol IP (As preservative)	0.945 % v/v
Water for Injections IP	q.s.

PREDNISOLONE crystalline injectable suspension was developed for use in the veterinary field. It has proved to be highly effective in the treatment of acetonemia in cattle (bovine ketosis). Following the injection of PREDNISOLONE crystalline Suspension, the blood glucose level is restored to normal in short time, appetite and milk yield improves, and circulating ketone bodies return to a normal level. It is three to five times more active than hydrocortisone, (natural cortizones) but does not affect mineral balance as that of the other potent corticosteroids.

INDICATIONS

PREDNISOLONE crystalline Suspension is used in treating ketosis in dairy cattle, rheumatoid arthritis, bursitis inflammatory and allergic conditions in large and small animals in conjunction with appropriate antibiotic or chemotherapeutic agent when indicated.

DOSE AND ADMINISTRATION

For systemic treatment, injection is given deep into the muscle, 2-4 times if necessary at an interval of 24 hours.

Cattle	: 50 - 100 - 200 mg
Calves	: 25 - 50 mg
Piglets, dogs, cats	: 10 - 30 mg

For local treatment injection is given in the part involved carefully under strict aseptic conditions. Contra indications include diabetes mellitus & osteoporosis. In such cases, crystalline suspension must not be administered more than twice.

Injection into Joints & tendon sheaths

Large animals : 25 - 75 mg or more

Dogs & Cats : 5 - 20 mg or more

Injection into Bursae

10 - 50 mg or Bursae

Subconjunctival injection

1.25 - 12.5 mg

Retrobulbar injection

2.5 - 12.5 mg

Infiltration (as in periostitis etc.)

10 - 50 mg

Injection into infected tissues are contra-indicated. The preparation should be shaken well before use. Use fine needle for local therapy.

STORAGE : Store protected from light at a temperature not exceeding 30°C

PRECAUTIONS : Shake well before use.

Keep out reach of children.

WITHDRAWAL PERIOD :

Milk - 3 days

Meat - Cattle, Sheep & Goat - 5 days & Pig - 28 days

PRESENTATION :

Vial of 10 ml

Marketed by:

Intervet India Pvt. Ltd.,
Bldg. B-1, Godown No. 12 to 16, and 12-A to 17-A,
Sagar Complex, Owali Village, Bhiwandi,
Thane - 421 302 (India)

Manufactured by: Akums Drugs & Pharmaceuticals Ltd.
2, 3, 4 & 5, Sector - 68, I.I.E., SIDCUL, Ranipur,
Haridwar- 249 403 (India)

