

Rx
Tetracycline Hydrochloride
Water Soluble Vet



NOT FOR HUMAN USE - FOR ANIMAL TREATMENT ONLY

SCHEDULE H PRESCRIPTION DRUG - CAUTION Not to be sold by retail without the prescription of a Registered Veterinary Practitioner only.

COMPOSITION

Each gram contains:

Tetracycline Hydrochloride IP 50 mg

Tetracycline hydrochloride belongs to the group of broad-spectrum antibiotics. Chemically it can be regarded as the active basis of the antibiotics, oxy- and chlortetracycline, which were discovered earlier. The antibiotic range of Tetracycline corresponds to that of oxy- and chlortetracycline and covers almost all gram-positive and gram-negative organisms. It has no action on *Proteus vulgaris* and certain strains of *Pseudomonas aeruginosa*.

Tetracycline Water Soluble Powder is used for the prevention and treatment of a wide variety of infections of the respiratory, gastrointestinal, genital and urinary tracts, in large and small animals and birds.

Tetracycline therapy offers many advantages:-

High degree of efficacy-Rapid results - Ease of administration-Ready solubility-Well-absorbed from the gastro-intestinal tract-Broad spectrum of activity.

INDICATIONS

Poultry : Infectious coryza, Fowl cholera, White Bacillary diarrhoea, Fowl typhoid, CRD, Psittacosis (Ornithosis), Spirochaetosis. To combat secondary infections

associated with viral diseases or to prevent infections during stress periods such as vaccination, extremes of temperature, transport etc.

Large and small animals:

Infections following operations, wounds, delivery, dystokia, retention of placenta, Haemorrhagic septicaemia, Anthrax, Black quarter, Calf pneumonia, Calf diphtheria, Scours, Joint ill, Strangles, Contagious caprine pleuro-pneumonia, Foot rot, Acute mastitis, Leptospirosis, Acute metritis, Pneumonia, To combat secondary infections associated with Rinderpest, Mucosal disease complex, Foot and Mouth Disease and Canine Distemper.

DOSE AND ADMINISTRATION :

Tetracycline Water Soluble powder can be given mixed in milk, liquid feed or drinking water, or as a drench or electrolyte.

Large animals : 2.5-5g / 15 kg body weight

Small animals : 1g / kg body weight

Poultry :

Prevention : 2.5 g in 4.5 litres of drinking water

Treatment : 5 g in 4.5 litres of drinking water

NOTE: The above dosage may be increased or doubled in case of very severe infections.

STORAGE:

Store protected from light and moisture.

PRESENTATION : 100 g sachet.

WITHDRAWAL PERIOD :

Milk - 7 days, Egg - 7 days

Meat - Cattle, Pig, Sheep & Goat: 28 days

Poultry : 5 days

Marketed by : **Intervet India Pvt. Ltd.**

Bldg. B-1, Godown No. 12 to 16, and 12-A to 17-A,
Sagar Complex, Owali Village, Bhiwandi, Thane - 421302, India.

Consumer Care Number: +91-20-66294700/01

E-mail: info@india@merck.com

Website: www.msdl-animal-health.co.in

Manufactured by :

EMIL Pharmaceutical Ind. Pvt. Ltd.
N-50, MIDC Tarapur, Boisar,
Dist. Palghar 401506, India.

