

FOR THE USE ONLY OF A VETERINARIAN OR A HOSPITAL OR A LABORATORY OR A FARM

Oxyclozanide Suspension I P

Tolzan® F VET Fluke Drench



NOT FOR HUMAN USE-FOR ANIMAL TREATMENT ONLY

Warning

To be sold by retail on the prescription of a Veterinary Doctor only.

COMPOSITION

Oxyclozanide IP 3.4% w/v
Excipients q.s.

INDICATIONS

For the treatment and control of fascioliasis, amphistomiasis in cattle, buffaloes, sheep and goats. Tolzan® F Vet suspension eliminates practically all flukes (Fascioliasis) present in the bile duct. Moneiz tapeworm segments are also removed. For immature tissue flukes of sheep, 3 times the therapeutic dose of Oxyclozanide suspension is highly effective. Tolzan® F Vet suspension is effective in the treatment of acute fascioliasis in sheep. Tolzan® F Vet suspension is also effective against para amphistome

SPECIAL FEATURES

Tolzan® F Vet suspension, when given to ewes 6 weeks before lambing, improves the subsequent performance of the lamb. Tolzan® F Vet suspension effectively controls liver flukes and amphistomiasis infection and increases productivity of the dairy animals.

DOSE AND ADMINISTRATION

Tolzan® F Vet suspension can be administered orally direct or through feed or in the form of suspension to each individual animal.

DOSE

For Tolzan® F Suspension : (Shake well before use)

Cattle & Buffalo :	10 mg of Oxyclozanide per kg b/w	
	30 ml per 100 kg b/w upto a maximum 100 ml	
Sheep & Goat :	15 mg / kg (Oxyclozanide)	
	Bodyweight	Dose
	Upto 15 kg	5 ml
	15 - 30 kg	10 ml
	30 - 45 kg	15 ml
	45 kg and over	20 ml

CONTRAINDICATIONS

No side effects have been observed in sheep. In cattle, Tolzan® F Vet suspension occasionally causes slight softening of the faeces with transient loss of appetite. Tolzan® F Vet suspension does not affect the quality of milk.

SAFETY

Tolzan® F Vet suspension is safe for debilitated, pregnant and lactating animals. Diet changes are not required and may be given at the same time as treatment for other conditions or during vaccination.

PRECAUTIONS

Due care must always be given to the physical condition of the animals receiving Tolzan® F Vet suspension those in advanced pregnancy or under stress, poor nutrition and handling etc. Animals should not be slaughtered for human consumption for a period of 14 days after treatment with Tolzan® F Vet suspension.

STORAGE

Store protected form light.
Shake well before use

PRESENTATION

Tolzan® F Vet suspension : 90 ml & 1 Ltr container

WITHDRAWAL PERIOD

Milk - 07 days
Meat - 14 days

* Registered Trademark

Marketed by : **Intervet India Pvt. Ltd.**

Bldg. B-1, Godown No. 10 to 16, and 12-A to 17-A,
Sagar Complex, Owali Village, Bhiwandi, Thane - 421302,
Maharashtra, India.

Consumer Care Number: +91-20-66294700/01

E-mail: infoin-india@merck.com

Website: www.msd-animal-health.co.in

Manufactured by :
EMIL Pharmaceutical Ind. Pvt. Ltd.
N-50, MIDC Tarapur, Boisar,
Dist. Palghar 401506,
Maharashtra, India.



MSD
Animal Health

PH08000223 55 X 105 mm